



Molecular phylogeny and biotechnological potential of bacterial endophytes associated with *Malpighia emarginata*

V. Specian¹, A.T. Costa¹, A.C. Felber¹, J.C. Polonio¹, J.L. Azevedo² and J.A. Pamphile¹

¹Departamento de Biotecnologia, Genética e Biologia Celular, Universidade Estadual de Maringá, Maringá, PR, Brasil

²Departamento de Genética, Escola Superior de Agricultura “Luiz de Queiroz”, Universidade de São Paulo, Piracicaba, SP, Brasil

Corresponding author: J.A. Pamphile
E-mail: prof.pamphile@gmail.com

Genet. Mol. Res. 15 (2): gmr.15027777

Received October 5, 2015

Accepted December 3, 2015

Published April 26, 2016

DOI <http://dx.doi.org/10.4238/gmr.15027777>

ABSTRACT. Acerola (*Malpighia emarginata*) is a shrub native to tropical and subtropical climates, which has great commercial interest due to the high vitamin C content of its fruit. However, there are no reports of the endophytic community of this plant species. The aim of this study was to verify the genetic diversity of the leaf endophytic bacterial community of two varieties (Olivier & Waldy Cati 30) of acerola, and to evaluate their biotechnological ability by assessing their *in vitro* control of pathogenic fungi and the enzymatic production of cellulase, xylanase, amylase, pectinase, protease, lipase, esterase, and chitinase. In total, 157 endophytic bacteria were isolated from the leaves of two varieties of the plant at 28° and 37°C. Phylogenetic analysis confirmed the molecular identification of 58 bacteria, 39.65% of which were identified at the species level. For the first time, the genus *Aureimonas* was highlighted as an endophytic bacterium. Furthermore, 12.82% of the isolates inhibited the growth of all phytopathogens evaluated and

at least one of the above-mentioned enzymes was produced by 64.70% of the endophytes, demonstrating that *M. emarginata* isolates have potential use in biotechnological studies.

Key words: Endophytes; Acerola; Phylogenetic analysis; Antagonistic activity; Enzymes

INTRODUCTION

Malpighia emarginata D.C., also referred to as West Indian cherry, Caribbean cherry, Barbados cherry, and popularly termed acerola in Brazil, is a wild plant that originated in southern Mexico and in the northern region of South America (Carrington and King, 2002). This species was introduced to Brazil about 65 years ago, and its production is concentrated mainly in the northeastern, southeastern, and southern regions of the country (Santos et al., 2012). Brazil is considered the biggest producer, consumer, and exporter of acerola in the world (Cavichioli et al., 2014). This fruit is now used widely for preparing juices and for general consumption because of its high levels of vitamin C (Cavichioli et al., 2014).

Endophytes are microorganisms that inhabit the interior of plant tissues during all or part of their life cycle, without causing any visible symptoms (Azevedo et al., 2000; Kusari et al., 2012). All plants harbor a diverse community of endophytic bacteria, which live in symbiosis with plants, passing through epiphytic and endophytic colonization. These bacteria may be isolated from the vegetal tissue after surface disinfection (Lacava and Azevedo, 2013).

These bacteria stimulate plant growth, yield, can act as biological control agents, and produce a variety of natural products that have potential use in medicine, agriculture, and industry. They may also remove contaminants from soil by promoting phytoremediation and may influence soil fertility via phosphate solubilization and nitrogen fixation (Anand et al., 2014).

Molecular approaches based on PCR amplification and sequencing of the 16S rRNA gene are important tools for use in taxonomy and bacterial identification, and are used to identify new pathogens and to define complex microbial communities (Schlaberg et al., 2012). The function of the 16S rRNA gene has not changed over the course of evolution, suggesting that random-sequence alterations are a relatively accurate measure of time.

As they colonize a similar ecological niche as phytopathogens, endophytic bacteria are suitable for use as biocontrol agents. Various reports have shown that endophytic microorganisms have the ability to control several plant pathogens (Chebotar et al., 2015). The “biocontrol activity” of microorganisms is defined as their ability to reduce populations of target species of antagonist-organisms through a variety of ecological mechanisms, including pathogenesis, competition within the ecological niche, and the production of compounds that inhibit their growth and development (Chebotar et al., 2015).

An efficient antagonist against several fungal pathogens is *Bacillus subtilis*, due to its production of antifungal compounds, antibiotics, and proteases, causing this species to be extensively used in agricultural systems (Soria et al., 2012). The inhibitory effect of live bacteria and their thermostable metabolites on the *in vitro* growth of *Fusarium circinatum*, the phytopathogen that induces cancer of the pine (*Pinus taeda* L.), were demonstrated by four endophyte isolates of *B. subtilis* and one isolate of *Burkholderia* sp. In this study, all bacteria displayed an antagonist effect on pathogen growth and the thermostable metabolites inhibited

fungal growth by over 50%, demonstrating that endophytic bacteria constitute an alternative method of *F. circinatum* control (Soria et al., 2012; Chebotar et al., 2015).

Microbial enzymes are of great importance because they act as biocatalysts in several industries, including biotechnology, agricultural, and pharmaceutical industries. Bacteria have been used in the production of enzymes for years, and microbial cultivated enzymes have replaced animal or plant enzymes (Duza and Mastan, 2013). Bacterial hydrolytic enzymes like cellulases and pectinases are also used by endophytic bacteria to penetrate plant tissues (Gujral et al., 2013). Due the high production capability, low cost, and susceptibility to genetic manipulation, enzymatic production processes have generated much biotechnological interest (Castro et al., 2014).

Prompted by the great potential of endophyte microorganisms and the lack of information on endophytes isolated from acerola leaves, the aim of this study was to verify the genetic diversity of the leaf bacterial endophyte community of two varieties of *M. emarginata* (Olivier & Waldy Cati 30), and to evaluate their biotechnological ability in relation to the production of enzymes and the *in vitro* control of phytopathogenic fungi.

MATERIAL AND METHODS

Leaf sampling

Healthy and undamaged adult leaves were randomly collected from two varieties (Olivier & Waldy Cati 30) of *M. emarginata* on the 170-ha Experimental Farm of the Universidade Estadual de Maringá, Iguatemi, in the northwestern region of the State of Paraná, Brazil, 23° 25' S and 51° 57' W, altitude 550 m. Leaves were collected under a mean temperature of 22.3°C and relative air humidity 60%, with mean rainfall index of 29.7 mm. Data on temperature and rainfall were retrieved from <http://www.inmet.gov.br> (National Institute of Meteorology), station A835. The leaf material was processed in the Laboratory of Microbial Biotechnology of the State University of Maringa, Maringá, Paraná, Brazil (BIOMIC-UEM).

Isolation and maintenance of endophytic bacteria

Leaves were washed in running water to remove residue, cleansed with 0.01% Tween 80 aqueous solution (Synth), and washed twice in autoclaved distilled water. Surface disinfection was undertaken under running water with immersion in 3% sodium hypochlorite (active chlorine) for 4 min. The efficiency of the method was evaluated by spreading 100 µL of the water used on Petri dishes containing LB culture medium (10 g/L tryptone, 5 g/L yeast extract, 10 g/L NaCl, 15 g/L agar, in 1 L distilled water; pH 7.0), supplemented with fungicide Benomyl (50 µg/mL in absolute ethanol). All procedures were performed separately for each variety.

Five leaf fragments (5 mm²) were placed on each Petri dish containing LB medium supplemented by Benomyl, and incubated at 28° and 37°C for 3-4 days until bacterial growth was evident on most fragments. Frequency of isolation (FI) was determined by $FI = (\text{number of fragments colonized bacteria} / \text{total number of leaf fragment}) \times 100$.

Isolates were transferred to plates containing LB medium and further incubated at 28° and 37°C, complying with the isolation temperature. After 2-5 days, depending on growth

requirements required for each isolate, bacteria were purified using the streaking technique to obtain pure colonies. The codes used to distinguish the origin of isolates and temperatures of isolation were according to the Table 1.

Table 1. Codification used to distinguish the origin of endophytes (plant variety) and temperature used to growth of the bacteria.

Codes	Description	Example 1	Example 2
First letter	Abbreviation of the group.	G	G
First number	Number attributed to group based on the morphological characteristics of the isolates.	1	2
Second letter	The variety from where the endophyte was retrieved, O (Olivier) and W (Waldy Cati 30).	O	W
Second number	Number given to the isolate within the group.	33	2
Letter between brackets	Temperature of isolation, A (37°C) and B (28°C).	(A)	(B)
Results		G1O33 (A)	G2W2 (B)

Previous assessments showed that bacterial isolates grew better in TSA culture medium (Tryptone Soy Agar, HIMEDIA) (data not shown). All isolates were kept in TSA and cultures of pure colonies were maintained at 28° and 37°C for 2 days prior to the start of each experiment. Isolates were preserved by freezing in 30% glycerol and were stored at -20°C.

Identification of endophyte isolates

Bacterial DNA was extracted following the method described by Nogueira et al. (2004), with modifications. Endophytic bacteria were grown in 5 mL LB broth for 24 h at 28°C. Next, 400 µL of the suspension was transferred to microtubes, to which 400 µL of saturated phenol solution was added. The mixture was stirred (Vortex®) and centrifuged at 16,000 g for 5 min. The supernatant (water phase) was transferred to another microtube and the phenolic stage replicated. After centrifugation, the supernatant was transferred to another microtube containing 400 µL chloroform. The microtube was vortexed and centrifuged for 5 min at 16,128 g. The water layer was transferred to another microtube to which 1 mL ethanol was added. The DNA extraction was completed when the microtube was centrifuged for 3 min at 16,128 g, and a DNA pellet was formed. The ethanol was disposed of and the tubes were exposed to 37°C for 30 min to allow any ethanol to evaporate. The pellet was then eluted in 15 µL autoclaved ultrapure water.

The 16S region of rDNA was amplified following the method described by Procópio et al. (2009), with modifications. The PCR was performed in a final volume of 50 µL containing 5 µL buffer (200 mM Tris-HCl, pH 8.4, 500 mM KCl), 5 µL dNTPs (2.5 mM), 3 µL of each primer (Invitrogen pmol/µL 10⁻¹) (R1378: 5'-CGGTGTGTACAAGGCCCGGAACG-3' and PO27F: 5'-GAGAGTTTGATCCTGGCTCAG-3'), 0.4 µL Taq DNA polymerase (5 U/µL), 3.75 µL MgCl₂ (50 mM), 27.85 µL ultrapure water, and 2 µL sample DNA (10-20 ng/µL). PCR conditions comprised initial denaturation at 94°C for 4 min, followed by 25 cycles: denaturation at 94°C for 30 s; annealing at 63°C for 1 min; extension at 72°C for 1 min, and a final extension at 72°C for 7 min. Amplification products were purified with two enzymes, shrimp alkaline phosphatase (SAP) and exonuclease I (EXO). Reactions were performed using 8 µL PCR product, 0.5 µL EXO (10 U/µL), and 1 µL SAP (1 U/µL), and incubated in a thermocycler for 1 h at 37°C, followed by 15 min at 80°C, and were conserved at 4°C. Samples were sequenced by ACTGene Análises Moleculares Ltd. (Ludwigbiotec).

Sequences were then analyzed and edited. Isolates were identified from the percentage identity and sequence coverage compared to those deposited in GenBank (<http://www.ncbi.nlm.nih.gov>) with BLASTn.

A similarity dendrogram was generated for phylogenetic analyses using the sequences obtained by sequencing and those deposited in GenBank. Sequences were paired by Clustal W and with the dendrogram produced by MEGA 6 (Tamura et al., 2013), by the neighbor-joining method, p-distance for nucleotides with the pairwise gap deletion and bootstrap with 10,000 replications.

All rDNA 16S gene sequences in the current study were analyzed with Decipher to verify the presence of chimeras. After identification, sequences were compared with those belonging to the type strain from the LPSN database (<http://www.bacterio.net>). Sequences were compared using BLASTn with the Entrez Query option (with the addition of the accession No. of type strain sequences). All rDNA 16S gene sequences in the current study were deposited in the NCBI GenBank under the accession Nos. KR005461 to KR005518 (Table 2).

Table 2. Identification of endophytic bacteria of *Malpighia emarginata* based on rDNA sequencing.

Endophyte strain (GenBank)	Plant variety	Isolation temp.	BLAST (GenBank)*		Type strain sequence**		Taxonomic identification***
			Accession No. with highest identity	Identity (%)	Accession No.	Identity (%)	
G1017 (A) (KR005461)	Olivier	37°C	KM042087.1	100	-	-	<i>Enterobacter</i> sp
G1033 (A) (KR005462)	Olivier	37°C	JN585689.1	96	-	-	<i>Staphylococcus</i> sp
G1038 (A) (KR005463)	Olivier	37°C	KM250109.1	98	-	-	<i>Staphylococcus</i> sp
G2023 (A) (KR005464)	Olivier	37°C	KM203879.1	99	-	-	<i>Staphylococcus</i> sp
G2039 (A) (KR005465)	Olivier	37°C	KM083802.1	99	-	-	<i>Staphylococcus</i> sp
G2043 (A) (KR005466)	Olivier	37°C	KM015449.1	97	AF094748	97%	<i>Pseudomonas stutzeri</i>
G3037 (A) (KR005467)	Olivier	37°C	KF017561.1	100	-	-	<i>Bacillus</i> sp
G3029 (A) (KR005468)	Olivier	37°C	KM083802.1	100	-	-	<i>Staphylococcus</i> sp
G5014 (A) (KR005469)	Olivier	37°C	KM042087.1	100	-	-	<i>Enterobacter</i> sp
G5025 (A) (KR005470)	Olivier	37°C	FJ842658.1	100	-	-	<i>Bacillus</i> sp
G5027 (A) (KR005471)	Olivier	37°C	EU071560.1	99	-	-	<i>Bacillus</i> sp
G5047 (A) (KR005472)	Olivier	37°C	KM052592.1	100	CP000002.3	99%	<i>Bacillus licheniformis</i>
G5048 (A) (KR005473)	Olivier	37°C	KM052592.1	99	CP000002.3	99%	<i>Bacillus licheniformis</i>
G5049 (A) (KR005474)	Olivier	37°C	JN812081.1	99	CP000002.3	99%	<i>Bacillus licheniformis</i>
G8031 (A) (KR005475)	Olivier	37°C	KM052592.1	96	CP000002.3	96%	<i>Bacillus licheniformis</i>
G8036 (A) (KR005476)	Olivier	37°C	KM093730.1	100	-	-	<i>Bacillus</i> sp
G8037 (A) (KR005477)	Olivier	37°C	KM203879.1	100	-	-	<i>Staphylococcus</i> sp
G8043 (A) (KR005478)	Olivier	37°C	KF017550.1	100	-	-	<i>Bacillus</i> sp
G8050 (A) (KR005479)	Olivier	37°C	KM016934.1	100	-	-	<i>Bacillus</i> sp
G1W28 (A) (KR005480)	Waldy Catí 30	37°C	KM093730.1	99	-	-	<i>Bacillus</i> sp
G4W14 (A) (KR005481)	Waldy Catí 30	37°C	KM042089.1	100	-	-	<i>Enterobacter</i> sp
G101 (B) (KR005482)	Olivier	28°C	HG810389.1	98	Z93440.1	97%	<i>Acinetobacter johnsonii</i>
G108 (B) (KR005483)	Olivier	28°C	KJ995847.1	99	Z93440.1	99%	<i>Acinetobacter johnsonii</i>
G109 (B) (KR005484)	Olivier	28°C	KJ803945.1	92	D84020	90%	<i>Pseudomonas putida</i>
G1024 (B) (KR005485)	Olivier	28°C	AM184286.1	95	D84020	92%	<i>Pseudomonas putida</i>
G1032 (B) (KR005486)	Olivier	28°C	KJ803945.1	99	D84020	96%	<i>Pseudomonas putida</i>
G1038 (B) (KR005487)	Olivier	28°C	KM250110.1	100	D16281	100%	<i>Bacillus thuringiensis</i>
G1039 (B) (KR005488)	Olivier	28°C	KJ880015.1	100	Z93440.1	99%	<i>Acinetobacter johnsonii</i>
G1047 (B) (KR005489)	Olivier	28°C	HQ941660.1	100	-	-	<i>Staphylococcus</i> sp
G2026 (B) (KR005490)	Olivier	28°C	KJ943979.1	99	-	-	<i>Bacillus</i> sp
G2029 (B) (KR005491)	Olivier	28°C	KM108311.1	100	D16281	99%	<i>Bacillus thuringiensis</i>
G2034 (B) (KR005492)	Olivier	28°C	KM250109.1	100	-	-	<i>Staphylococcus</i> sp
G302 (B) (KR005493)	Olivier	28°C	HQ880685.1	85	-	-	<i>Bacillus</i> sp
G306 (B) (KR005494)	Olivier	28°C	KJ803945.1	100	D84020	97%	<i>Pseudomonas putida</i>
G3014 (B) (KR005495)	Olivier	28°C	KM042089.1	100	-	-	<i>Enterobacter</i> sp
G3015 (B) (KR005496)	Olivier	28°C	KM250110.1	100	D16281	100%	<i>Bacillus thuringiensis</i>
G3025 (B) (KR005497)	Olivier	28°C	KJ880015.1	100	Z93440.1	99%	<i>Acinetobacter johnsonii</i>
G3046 (B) (KR005498)	Olivier	28°C	HMS67114.1	99	D16281	98%	<i>Bacillus thuringiensis</i>
			KM108311.1	99	-	-	
G404 (B) (KR005499)	Olivier	28°C	KF018921.2	99	-	-	<i>Bacillus</i> sp
G5018 (B) (KR005500)	Olivier	28°C	HE681737.1	98	X77445	97%	<i>Microbacterium testaceum</i>
G7036 (B) (KR005501)	Olivier	28°C	KF018921.2	100	-	-	<i>Bacillus</i> sp
G8043 (B) (KR005502)	Olivier	28°C	KF018921.2	100	-	-	<i>Bacillus</i> sp
G8049 (B) (KR005503)	Olivier	28°C	KM186185.1	100	-	-	<i>Bacillus</i> sp
G8051 (B) (KR005504)	Olivier	28°C	KF891394.1	89	X82598	88%	<i>Cellulomonas hominis</i>
			Q660180.1	89	-	-	
G1W1 (B) (KR005505)	Waldy Catí 30	28°C	KJ744023.1	99	AJ536198	99%	<i>Micrococcus luteus</i>

*Percent identity based on the comparison of rDNA sequences of endophytic bacteria deposited in GenBank using BLAST. **Comparison of endophytic bacteria with bacteria from the LPSN database (List of Prokaryotic names with Standing in Nomenclature) (<http://www.bacterio.net/>) and compared with BLAST. ***Taxonomic identity of endophytic bacteria based on the following analyses: BLAST (*), type strain sequences (**), and phylogenetic analysis.

Antagonistic activity of endophytic isolates against phytopathogenic fungi *in vitro*

Isolates of endophytic bacteria from the leaf tissue of *M. emarginata* were used to perform an antagonism assay against seven phytopathogens (Table 3).

Table 3. Phytopathogenic fungi identified in an antagonism assay.

Species	Disease	Host plant	Laboratory
<i>Glomerella</i> sp (CNPUV 102)	Grape rotteness, anthracnose	Grapevine, acerola	EMBRAPA Grape and Wine of Bento Gonçalves - RS, Brazil
<i>Sclerotinia sclerotiorum</i>	White Mold; white rotteness	Soy bean, sun flower, canola, peas, beans, potato	Laboratory of Genetics of Microorganisms "João Lúcio de Azevedo" ESALQ - USP, SP, Brazil
<i>Monilophthora perniciosa</i>	Witches' broom	Cocoa	Laboratory of Genetics of Microorganisms "João Lúcio de Azevedo" ESALQ - USP, SP, Brazil
<i>Fusarium solani</i>	Red rotteness of the root	Soy bean	Laboratory of Genetics of Microorganisms "João Lúcio de Azevedo" ESALQ - USP, SP, Brazil
<i>Didymella bryoniae</i>	Gummy stem blight	Watermelon, melon	Laboratory of Genetics of Microorganisms "João Lúcio de Azevedo" ESALQ - USP, SP, Brazil
<i>Alternaria</i> sp (CNPUV 674)	Early blight; black spot	Acerola, mango, tomato, potato, eggplant	EMBRAPA Grape and Wine of Bento Gonçalves - RS, Brazil
<i>Fusarium oxysporum</i> (ATCC 2163)	Fusariosis	Acerola	Institution André Tosello, Campinas, SP, Brazil

In vitro antagonism assays were conducted by dual culture to verify the capacity of the endophytes to inhibit phytopathogen growth. Fungi were previously grown on potato dextrose agar (PDA) culture medium at 28°C for 7 days. The 39 endophytic bacteria were grown in TSA medium at 28°C for 24 h. After this period, each bacterium was striated along two edges of a Petri dish (1 cm from the margin) containing PDA. After 48 h, each fungus (6-mm discs) was transferred to the center of the dish (3 cm from each bacterial striate). Two controls were used: in the first control, phytopathogens were inoculated in the middle of the Petri dish to assess their maximum growth; in the second control, 6-mm discs of the same phytopathogens were placed at the edges of the dishes equidistant at 4 cm. Tests were performed in triplicate and all dishes were incubated at 28°C for 6 days. The capacity of isolates to inhibit pathogen fungi was determined by measuring the growth size (in cm) of the pathogen in the experiment and in the controls.

Evaluation of the enzymatic activity of endophytic isolates

M. emarginata isolates were tested for their capacity to produce cellulase, xylanase, amylase, pectinase, protease, lipase, esterase, and chitinase in solid culture media. Thirty-four strains of endophytic bacteria were grown in tryptone soy broth (TSB, HIMEDIA) liquid medium for 48 h at 28°C. The bacteria were then stirred and eluted in 5 mL 1% saline solution to obtain a concentration of 10⁸ colony-forming units per milliliter (CFU/mL), equivalent to 0.5 on the McFarland scale. Next, 2 µL of the culture was inoculated on Petri dishes containing minimal medium for bacteria, -M9 200 mL/L stock solution (64 g/L Na₂HPO₄·7H₂O; 15 g/L KH₂PO₄; 2.5 g/L NaCl; 5 g/L NH₄Cl; 2.0 mL/L 1 M MgSO₄; 10 g/L carbon source; 0.1 mL/L CaCl₂ 1 M; 15 g/L agar, pH 7.2, with the following carbon sources [amylase 0.5%, yeast extract, and 1% soluble starch (v/v)] cellulase [0.5% yeast extract and 1% carboxymethylcellulose (v/v)], pectinase [0.5% yeast extract and 1% pectin (v/v), pH 8.0], xylanase [0.5% yeast extract and 1% xylan (v/v)]. To evaluate lipolytic and esteratic activity, lipase/esterase medium was used (10 g/L peptone; 5 g/L NaCl; 0.1 g/L CaCl₂·H₂O; 15 g/L agar; pH 7.4) with 1% (v/v) Tween 20 for lipase and Tween 80 for esterase. To investigate proteolytic activity, protease medium was used (5 g/L tryptone; 2.5 g/L yeast extract; 1.0 g/L glucose; 2.5 g/L NaCl; 15 g/L agar; pH 7.0); after autoclaving, 100

mL skimmed milk was added per 900 mL medium. To detect enzymatic activity, the incubation period ranged from 24 to 168 h at 28°C. To visualize the enzyme activities of cellulase, amylase, pectin, and protease, congo red dye, iodine tincture, HCl 5 N, and acetic acid were used, respectively; revealers were not required for the others, since enzymatic production could be visualized as a bright halo around the colonies (Oliveira et al., 2006). Commercial enzymes were used as a positive control; these included protease (*Aspergillus oryzae*), porcine pancreatic α -amylase, and pectinase (*Aspergillus niger*; Sigma). The negative control comprised 1% saline solution. Semi-quantitative enzymatic activity was evaluated by calculating the enzymatic index (EI) using the formula $EI = \text{halo diameter (cm)} / \text{diameter of the colony (cm)}$.

Statistical analyses

Experiments were randomized and statistically analyzed by analysis of variance (ANOVA); means were compared by the Scott-Knott test ($P < 0.05$) using Sisvar 4.3 (Ferreira, 2011).

RESULTS

Isolation and identification of endophytic bacteria

A total of 250 leaf fragments were sampled for each variety of *M. emarginata* and for each temperature; a total of 157 bacterial isolates were subdivided as follows: Olivier variety = 53 isolates (FI = 21.2%) obtained at 28°C; 50 isolates (FI = 20%) at 37°C; Waldy Cati variety 30 with 26 isolates (FI = 10.4%) obtained at 28°C and 28 isolates (FI = 11.2%) at 37°C.

These 157 endophytic bacteria were isolated and purified by the streaking method and grouped in morphogroups according to the morphological characteristics of color, colony formation, viscosity, development, and growth on TSA. In the case of the Olivier variety, bacteria isolated at 28°C were placed in nine groups and those isolated at 37°C were placed in a further eight groups. For the Waldy Cati variety, 30 bacteria were isolated at 28°C and placed in six groups, and those isolated at 37°C were also grouped into six groups. In total, 63/157 bacteria were randomly selected and used for further analyses.

Sequencing analyses of 16S rDNA led to the identification of 58 isolates distributed in nine genera: *Bacillus* (corresponding at 36.20% of the isolates), *Staphylococcus* (17.24%), *Enterobacter* (12.06%), *Pseudomonas* and *Microbacterium* (10.34% each), and *Acinetobacter* (6.89%). The genera *Aureimonas*, *Cellulomonas*, and *Micrococcus* were represented by a single endophyte. The endophyte sequences of *M. emarginata* shared 85-100% identity with those available in GenBank.

Molecular phylogeny analysis of endophytic bacteria of *M. emarginata* based on rDNA sequencing

The molecular phylogeny analysis divided the endophytic bacteria into 10 main clades (Figure 1).




Figure 1. Phylogenetic tree of endophytic bacteria isolated from *Malpighia emarginata* D.C. with other bacteria deposited in GenBank, built by the neighbor-joining grouping method with the p-distance for nucleotides and the pairwise gap-deletion parameter. Numbers on the tree represent the percentage of times the right hand group occurred on the same knot during the consensus assessment (bootstrap with 10,000 replications).

The genera *Bacillus* was grouped into Clade A, with 99% bootstrap probability (BP), where isolates G8043 (A), G8043 (B), and G404 (B) [100% identity with *Bacillus amyloliquefaciens* (KF018921.2)], isolate G8049 (B) [100% identity with *Bacillus methylotrophicus* (KM186185.1)], by BLAST analysis, and G7036 (B) [100% identified with *B. amyloliquefaciens* (KF018921.2)] were grouped with 100% BP with sequences deposited in GenBank, featuring different species of *Bacillus*. The latter were classified by molecular taxonomy as *Bacillus* sp. Isolate G5047 (A) [100% identity with *Bacillus licheniformis*

(KM052592.1)] was grouped with *B. licheniformis* with 69% BP, and was identified as *B. licheniformis*. Isolates G5O48 (A) [99% identity with *B. licheniformis* (KM052592.1)] by BLAST analysis, G5O49 (A) [99% identity with *Bacillus* sp (JM812081.1)], and isolate G8O31 (A) [96% identity with *B. licheniformis* (KM052592.1)] were grouped with 100% BP with those of isolates *B. licheniformis* in GenBank, and one isolate of *Bacillus* sp with 100% BP. These endophytic isolates were *B. licheniformis*. Isolate G1W28 (A) [99% identity with *Bacillus* sp (KM093730.1)], G5O25 (A) [100% identity with *Bacillus* sp (FJ842658.1)], G5O27 (A) [99% identity with *Bacillus pumilus* (EU071560.1)], G3O37 (A) [100% identified with *B. altitudines* (KF017561.1)], G8O50 (A) [100% identified with *B. pumilus* (KM016935.1)], and G8O36 (A) [100% identified with *Bacillus* sp (KM093730.1)] were grouped with several *Bacillus* species with 100% BP. These isolates should be taxonomically identified as *Bacillus* sp.

Also in Clade A, isolates G3O2 (B) [85% identified with *Bacillus* sp (HG880685.1)], G2O26 (B) [99% identity with *Bacillus* sp (KJ943979.1)], G1O38 (B) [100% identity with *Bacillus thuringiensis* (KM250110.1)], G3O15 (B) [100% identity with *B. thuringiensis* (KM250110.1)], G3O46 (B) [99% identified with *Bacillus* sp (HM567114.1) and *B. thuringiensis* (KM108311.1)], and G2O29 (B) [100% identity with *B. thuringiensis* (KM108311.1)] were grouped with all *B. thuringiensis* strains deposited in GenBank, as well as some identified as *Bacillus* sp, with 99% BP. Consequently, these strains may be taxonomically identified as *B. thuringiensis*. In Clade B, isolate G2W4 (B) [85% identity with *Bacillus circulans* (KF77099.1)] was grouped with the two isolates of *B. circulans* with 99% BP, and was classified as *B. circulans*. In Clade C, the isolate G3W20 (B) [93% identity with *Microbacterium* sp (JQ660075.1)] was placed in a sub-group with several strains of *Microbacterium* sp with 97% BP, corroborating its identification at a genus level. Isolate G5O18 (B) [98% identity with *Microbacterium testaceum* (HE681737.1)] was placed in a sub-group with other strains of *M. testaceum* with 89% BP, corroborating its identification at a species level. Isolate G3W25 (B) [99% identity with *Microbacterium* sp (FJ654469.1)] was placed in a sub-group with *Microbacterium* sp, corroborating its identification at a genus level. Isolate G2W7 (B) [98% identified with *Microbacterium paraoxydans* (LN556395.1)] was placed in a sub-group with other sequences of *M. paraoxydans* with 99% BP, corroborating its identification at a species level. The isolates G5W17 (B) and G2W6 (B) [99% identified with *Microbacterium* sp (EU104732.1)] by BLAST analysis, were placed in a sub-group with different species of the genus *Microbacterium*, corroborating the molecular taxonomic identification at a genus level.

In Clade D, isolate G8O51 (B) [89% identity with *Cellulomonas* sp (KF891345.1) and *Cellulomonas hominis* (JQ660180.1)] was placed in a sub-group with *C. hominis* with 99% BP, corroborating its identification at a species level (*C. hominis*).

In Clade E, isolate G1W1 (B) [99% identity with *Micrococcus* sp (KJ744023.1) and *Micrococcus luteus* (KJ733861.1)] was placed in a sub-group with 100% BP, corroborating identification at species level as *M. luteus*.

In Clade F, isolate G1W13 (B) [100% identity with *Aureimonas* sp (KJ685860.1) and 100% identity with *Aureimonas frigidaquae* (NR044195.1)] was placed in a sub-group with isolates that had the greatest identity by BLAST analysis, at 100% BP, confirming the identification of this species as *A. frigidaquae*.

In Clade G, isolates G2W14 (B), G2W15 (B), G5O14 (A), G1O17 (A), and G2W2 (B) [99, 100, 100, 100, and 99% identity with *Enterobacter* sp (KM042087.1), respectively],

and isolates G4W14 (A) and G3O14 (B) [100 and 100% identity with *Enterobacter* sp (KM042089.1), respectively] formed a sub-group with *Enterobacter* sp, with 100% BP, confirmed their identification at a genus level as *Enterobacter* sp.

In Clade H, endophytic isolates G3O6 (B) and G1O32 (B) [100 and 99% identity with *Pseudomonas putida* (KJ803945.1), respectively], formed a sub-group with the *P. putida* subclade with 62% BP, corroborating their identity at a species level. Isolates G2O43 (A) and G6W23 (B) [97 and 100% identity with *Pseudomonas stutzeri* (KM015449.1), respectively] formed a sub-group with *P. stutzeri* strains with 100% BP, corroborating their identity at a species level. Although, the isolates G109 (B) [92% *Pseudomonas putida* (KJ803945.1)] and G1O24 (B) [95% identity with *P. putida* (AM184286.1)], formed a sub-group with 99% BP, and with other species of *Pseudomonas* with 100% BP, they were identified as *Pseudomonas* sp.

In Clade I, isolates G1O1 (B) [98% identity with *Acinetobacter johnsonii* (HG810389.1)], G1O8 (B) [99% identity with *A. johnsonii* (KJ995847.1)], and endophytes G1O39 (B) and G3O25 (B) [both 100% identity with *A. johnsonii* (KJ880015.1)] were grouped in Clade H, which contains all sequences of *A. johnsonii*, confirming the taxonomic identity of the bacterial endophyte isolates at a species level.

In Clade J, the endophytic isolate G1O47 (B) [100% identity with *Staphylococcus haemolyticus* (HG941660.1)] grouped with *S. haemolyticus* with 100% BP, and was taxonomically identified as *S. haemolyticus*. Isolates G2W5 (B), G2O34 (B), and G1O38 (A) (B) [100, 100, and 100% identity, respectively, with *Staphylococcus warneri* (KM250109.1)], G8O37 (A) and G2O23 (A) [100 and 99% identity, respectively, with *Staphylococcus pasteurii* (KM203879.1)], G1O33 (A) [99% identity with *S. epidermidis* (JN585689.1)], G2O39 (A), G3W12 (B), and G3O29 (A) [99, 100, and 100% identity, respectively, with *Staphylococcus* sp (KM083802.1)] grouped with other isolates of the genus *Staphylococcus* sp with 54% BP. The molecular identification of these endophytic isolates is therefore *Staphylococcus* sp.

Assessment of antagonistic activity of endophytic bacteria toward phytopathogenic fungi

Although the 39 endophytic bacteria analyzed showed different levels of mycelium growth inhibition against the seven phytopathogens (Table 4), only three endophytic isolates, G4O4 (B) *Bacillus amyloliquefaciens*, G1W1 (B) *Micrococcus* sp, and G2W14 (B) *Enterobacter* sp, failed to show any antagonistic activity against any pathogen. Analyses of variance provided distinct groups for the pathogens analyzed for each endophyte (Table 5).

ANOVA revealed that the phytopathogen *Glomerella* sp had an antagonism index of 0-69.81% (Tables 4 and 5), with mean growth of the control being 8.48 cm. Based on variation in fungal growth, four groups of isolated endophytic bacteria with inhibition activity (IA) could be statistically distinguished. Isolate G8O43 (A) *Bacillus* sp (IA = 69.81%) was the best antagonist for *Glomerella* sp.

In the case of *S. sclerotiorum*, three statistically different groups were generated with antagonist indices ranging between 0 and 61.33%, with special reference to antagonism by isolate G1W28 (A) *Bacillus* sp (IA = 61.33%) (Figure 2).

Table 4. Antagonistic effect of endophytic bacteria on the growth of phytopathogenic fungi.

Endophytic bacteria	Antagonism index (%)*						
	GS**	SS	MP	DB	FS	AS	FO
G1O33 (A) <i>Staphylococcus</i> sp	23 ^c	0 ^c	17.22 ^b	-***	34.70 ^c	30.48 ^d	24.24 ^d
G1O38 (A) <i>Staphylococcus</i> sp	29.89 ^c	0 ^c	21 ^b	53 ^a	37.45 ^b	29.89 ^d	30.30 ^d
G2O23 (A) <i>Staphylococcus</i> sp	25.58 ^c	0 ^c	14.33 ^b	52.61 ^a	27.84 ^c	22.50 ^d	17.21 ^c
G2O39 (A) <i>Staphylococcus</i> sp	26.29 ^c	0 ^c	-	43.47 ^b	32.58 ^c	29.07 ^d	24.12 ^d
G2O43 (A) <i>Pseudomonas stutzeri</i>	25.70 ^c	0 ^c	23.55 ^b	51.04 ^a	31.58 ^c	22.50 ^d	23.27 ^d
G3O29 (A) <i>Staphylococcus</i> sp	24.41 ^c	0 ^c	21.33 ^b	47.65 ^a	28.71 ^c	24.97 ^d	29.09 ^d
G5O14 (A) <i>Enterobacter</i> sp	32.78 ^c	21 ^b	32.44 ^a	54.56 ^a	30.33 ^c	30.71 ^d	29.93 ^d
G5O27 (A) <i>Bacillus</i> sp	68.98 ^a	27.77 ^b	49.88 ^a	-	48.43 ^b	51.81 ^c	56.96 ^b
G5O47 (A) <i>Bacillus licheniformis</i>	32.19 ^c	0 ^c	27.22 ^b	-	26.84 ^c	27.19 ^d	25.93 ^d
G5O48 (A) <i>Bacillus licheniformis</i>	32.19 ^c	0 ^c	15.44 ^b	51.95 ^a	31.33 ^c	40.09 ^d	25.09 ^d
G5O49 (A) <i>Bacillus licheniformis</i>	24.37 ^c	0 ^c	24.33 ^b	37.59 ^b	34.08 ^c	25.43 ^d	26.66 ^d
G8O31 (A) <i>Bacillus licheniformis</i>	33.37 ^c	16.11 ^b	37.11 ^a	51.04 ^a	28.21 ^c	30.71 ^d	29.93 ^d
G8O36 (A) <i>Bacillus</i> sp	36.32 ^c	12.44 ^c	59.33 ^a	-	34.95 ^c	47.71 ^c	34.54 ^d
G8O37 (A) <i>Staphylococcus</i> sp	25.11 ^c	0 ^c	19.11 ^b	21.01 ^c	30.08 ^c	27.54 ^d	29.57 ^d
G8O43 (A) <i>Bacillus</i> sp	69.81 ^a	25 ^b	57.77 ^a	-	69.29 ^a	89.09 ^a	70.54 ^a
G8O50 (A) <i>Bacillus</i> sp	48.34 ^b	37.11 ^b	45.55 ^a	61.74 ^a	35.58 ^c	40.44 ^d	39.39 ^c
G1W28 (A) <i>Bacillus</i> sp	49.88 ^b	61.33 ^a	43.88 ^a	-	39.45 ^b	66.58 ^b	54.18 ^b
G4W14 (A) <i>Enterobacter</i> sp	50.94 ^b	23.77 ^b	28.33 ^b	58.22 ^a	39.45 ^b	37.74 ^d	41.09 ^c
G1O39 (B) <i>Acinetobacter johnsonii</i>	27.71 ^c	0 ^c	53.77 ^a	-	37.57 ^b	36.57 ^d	28.12 ^d
G1O47 (B) <i>Staphylococcus</i> sp	26.27 ^c	0 ^c	53.77 ^a	-	37.57 ^b	36.57 ^d	51.51 ^b
G2O34 (B) <i>Staphylococcus</i> sp	10.96 ^d	0 ^c	29.66 ^b	63.05 ^a	19.35 ^d	37.51 ^d	42.42 ^c
G3O6 (B) <i>Pseudomonas putida</i>	10.61 ^d	0 ^c	1.88 ^b	-	20.59 ^d	29.66 ^d	34.18 ^d
G3O14 (B) <i>Enterobacter</i> sp	26.17 ^c	0 ^c	15.55 ^b	-	29.71 ^c	23.79 ^d	32.36 ^d
G3O15 (B) <i>Bacillus thuringiensis</i>	33.72 ^c	0 ^c	25.44 ^b	56.78 ^a	29.96 ^c	25.55 ^d	19.27 ^e
G3O25 (B) <i>Acinetobacter johnsonii</i>	44.45 ^b	0 ^c	47.77 ^a	-	20.97 ^d	21.92 ^d	26.30 ^d
G4O4 (B) <i>Bacillus</i> sp	-	-	-	-	-	-	-
G5O18 (B) <i>Microbacterium testaceum</i>	26.29 ^c	30.55 ^b	13.88 ^b	-	25.34 ^c	24.85 ^d	19.87 ^c
G8O51 (B) <i>Cellulomonas hominis</i>	27.94 ^c	0 ^c	0 ^b	-	26.59 ^c	17.58 ^d	11.15 ^e
G1W1 (B) <i>Micrococcus luteus</i>	-	-	-	-	-	-	-
G1W13 (B) <i>Aureimonas frigidiquae</i>	13.32 ^d	0 ^c	12.11 ^b	36.94 ^b	16.85 ^d	18.17 ^d	6.06 ^e
G2W2 (B) <i>Enterobacter</i> sp	30.42 ^c	24.66 ^b	35.22 ^a	-	29.71 ^c	26.14 ^d	34.54 ^d
G2W4 (B) <i>Bacillus circulans</i>	30.66 ^c	0 ^c	-	-	24.96 ^c	21.10 ^d	14.18 ^e
G2W5 (B) <i>Staphylococcus</i> sp	34.19 ^c	0 ^c	46.00 ^a	-	47.81 ^b	46.89 ^c	41.61 ^c
G2W6 (B) <i>Microbacterium</i> sp	26.88 ^c	0 ^c	35.44 ^a	56.26 ^a	30.33 ^c	22.50 ^d	13.81 ^e
G2W14 (B) <i>Enterobacter</i> sp	-	-	-	-	-	-	-
G2W15 (B) <i>Enterobacter</i> sp	68.16 ^a	34.33 ^b	31.11 ^b	52.21 ^a	43.82 ^b	47.12 ^c	43.51 ^c
G3W25 (B) <i>Microbacterium</i> sp	43.98 ^b	0 ^c	18.22 ^b	-	24.46 ^c	20.75 ^d	14.78 ^e
G5W17 (B) <i>Microbacterium</i> sp	51.06 ^b	0 ^c	8.33 ^b	33.28 ^b	34.33 ^c	13.24 ^d	30.78 ^d
G6W23 (B) <i>Pseudomonas stutzeri</i>	35.96 ^c	0 ^c	13.00 ^b	39.16 ^b	48.06 ^b	20.51 ^d	27.75 ^d
Control 1****	0 ^d	0 ^c	0 ^b	0 ^d	0 ^c	0 ^d	0 ^c
Control 2*****	0 ^d	0 ^c	0 ^b	0 ^d	0 ^c	0 ^d	0 ^c

*Means of triplicate experiments followed by the same letter in the column did not differ by Scott-Knott's test ($P < 0.05$). **Phytopathogenic fungi: GS: Glomerella sp; SS: Sclerotinia sclerotiorum; MP: Moniliophthora perniciosa; DB: Didymella bryoniae; FS: Fusarium solani; AS: Alternaria sp; FO: Fusarium oxysporum. ***No antagonism. ****Control 1 = Petri plate with phytopathogens only (growth – diameter in cm, equivalent to 0% antagonism). *****Control 2 = Petri plate with two 6-mm discs of phytopathogens inoculated on the edges, equidistant at 4 cm (growth – diameter in cm, equivalent to 0% antagonism).

The *M. perniciosa* growth reduction by endophytes presents results in two statistically distinct groups of antagonism index. The isolate G8O43 (A) *Bacillus* sp had the greatest IA (57.77%).

Of note, the pathogen *Didymella bryoniae* had the least number of antagonists (48.71%) of the 39 bacterial isolates. Antagonism indices varied between 21.01 and 63.05%. Isolate G2O34 (B) *Staphylococcus* sp (IA = 63.05%) was the most effective antagonist.

To *Fusarium solani*, five different statistical groups of IA and four groups to *Alternaria* sp were obtained. Antagonism indexes for *F. solani* ranged between 16.85 and 69.29%, whereas for *Alternaria* sp, indices ranged between 0 and 89.09%. The endophytic isolate G8O43 (A) *Bacillus* sp was the most effective antagonist against the two phytopathogens.

The antagonism index of endophytic bacteria against *Fusarium oxysporum*, the phytopathogen that causes fusariosis in acerola, ranged from 0 to 70.54%, with the highest

Table 5. Antagonistic effect of endophytic bacteria on the growth of phytopathogenic fungi.

Phytopathogens	Grouping	Antagonism index (%)*	No. of endophytes
<i>Glomerella</i> sp	A	69.81-68.16	3
	B	51.06-43.98	6
	C	36.32-23.00	24
	D**	13.32-0	3
<i>Sclerotinia sclerotiorum</i>	A	61.33	1
	B	37.11-16.11	9
	C**	12.44-0	26
<i>Moniliophthora perniciosa</i>	A	59.33-32.44	13
	B**	31.11-0	22
<i>Dydmella bryoniae</i>	A	63.05-47.65	13
	B	43.47-33.28	5
	C	21.01	1
	D**	0	-
<i>Fusarium solani</i>	A	69.29	1
	B	48.43-37.45	9
	C	35.58-24.46	22
	D	20.97-16.85	4
	E**	0	-
<i>Alternaria</i> sp	A	89.09	1
	B	66.58	1
	C	51.81-46.89	4
	D**	40.44-0	30
<i>Fusarium oxysporum</i>	A	70.54	1
	B	56.96-51.51	3
	C	43.51-39.39	5
	D	34.54-23.27	19
	E**	19.87-0	8

*Means of triplicate experiments compared by the Scott-Knott test ($P < 0.05$). **Controls are also in this group.




Figure 2. Antagonistic activity of endophytic bacterial isolated from *Malpighia emarginata* against phytopathogens. **a.** phytopathogen *Sclerotinia sclerotiorum*; **b.** *Fusarium oxysporum*; **c.** *Alternaria* sp; **d.** antagonism mediated by isolate G1W28 (A) (*Bacillus* sp) against *S. sclerotiorum*; **e.** antagonism mediated by isolate G8O43 (A) *Bacillus* sp against *F. oxysporum*; **f.** antagonism mediated by isolate G8O43 (A) (*Bacillus* sp) against *Alternaria* sp.

indices obtained by the four isolates: G8O43 (A) *Bacillus* sp with (IA = 70.54%); G5O27 (A) *Bacillus* sp; G1W28 (A) *Bacillus* sp; and G1O47 (B) *Staphylococcus* sp. Variation among these isolates ranged between 51.51 and 56.96% in group B.

Enzymatic activity of endophytic isolates

Thirty-four endophytic bacteria isolated from *M. emarginata* were assessed for their enzymatic activity with regard to the production of protease, amylase, pectinase, lipase, esterase, cellulase, and xylanase. Protease accounted for the highest percentage (44.11%) of enzyme produced by *M. emarginata* endophytic bacteria isolates; followed by lipase (38.23%), amylase and pectinase (26.47%), and esterase (14.70%), whereas cellulase and xylanase had the lowest (11.76%) (Table 6).

Table 6. Protease, amylase, pectinase, lipase, esterase, cellulase, and xylanase activity of endophytic bacteria from *Malpighia emarginata*, reported as enzymatic index (EI) \pm standard deviation.

Endophytic bacteria	Enzymatic index (EI)*						
	Protease	Amylase	Pectinase	Lipase	Esterase	Cellulase	Xylanase
G2O23 (A) <i>Staphylococcus</i> sp	3.56 \pm 0.41 ^b	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G2O39 (A) <i>Staphylococcus</i> sp	4.80 \pm 0.33 ^a	1.25 \pm 0.12 ^c	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G2O43 (A) <i>Pseudomonas stutzeri</i>	0.00 \pm 0.00 ^e	2.15 \pm 0.35 ^b	2.30 \pm 0.34 ^b	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G3O29 (A) <i>Staphylococcus</i> sp	2.96 \pm 0.18 ^c	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G5O14 (A) <i>Enterobacter</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G5O27 (A) <i>Bacillus</i> sp	3.83 \pm 0.37 ^b	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	1.55 \pm 0.15 ^c	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G5O47 (A) <i>Bacillus licheniformis</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	1.32 \pm 0.19 ^c	1.72 \pm 0.19 ^b	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G8O37 (A) <i>Staphylococcus</i> sp	2.97 \pm 0.4 ^c	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G8O43 (A) <i>Bacillus</i> sp	2.22 \pm 0.45 ^d	1.34 \pm 0.20 ^c	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G8O50 (A) <i>Bacillus</i> sp	3.42 \pm 0.39 ^b	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	1.49 \pm 0.11 ^d	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1W28 (A) <i>Bacillus</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	1.24 \pm 0.09 ^c	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G4W14 (A) <i>Enterobacter</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O9 (B) <i>Pseudomonas putida</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O24 (B) <i>Pseudomonas putida</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O32 (B) <i>Pseudomonas putida</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O38 (B) <i>Bacillus thuringiensis</i>	2.17 \pm 0.35 ^d	1.65 \pm 0.15 ^c	2.21 \pm 0.14 ^b	1.57 \pm 0.26 ^d	1.23 \pm 0.06 ^b	1.41 \pm 0.08 ^b	0.00 \pm 0.00 ^f
G1O39 (B) <i>Acinetobacter johnsonii</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	2.03 \pm 0.16 ^a	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O47 (B) <i>Staphylococcus</i> sp	4.39 \pm 1.19 ^a	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G2O26 (B) <i>Bacillus subtilis</i>	2.69 \pm 0.41 ^c	1.73 \pm 0.15 ^c	2.32 \pm 0.24 ^b	1.73 \pm 0.16 ^b	1.21 \pm 0.03 ^b	1.41 \pm 0.09 ^b	1.53 \pm 0.16 ^c
G3O2 (B) <i>Bacillus</i> sp	2.15 \pm 0.21 ^d	1.77 \pm 0.11 ^c	2.26 \pm 0.31 ^b	1.35 \pm 0.06 ^c	0.00 \pm 0.00 ^e	1.39 \pm 0.07 ^b	1.52 \pm 0.14 ^c
G3O6 (B) <i>Pseudomonas putida</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G3O25 (B) <i>Acinetobacter johnsonii</i>	0.00 \pm 0.00 ^e	1.29 \pm 0.05 ^c	0.00 \pm 0.00 ^e	2.14 \pm 0.16 ^a	1.62 \pm 0.13 ^a	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G5O18 (B) <i>Microbacterium testaceum</i>	3.01 \pm 0.24 ^c	0.00 \pm 0.00 ^f	1.65 \pm 0.15 ^d	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	1.94 \pm 0.10 ^a	0.00 \pm 0.00 ^f
G7O36 (B) <i>Bacillus</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1O8 (B) <i>Acinetobacter johnsonii</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	1.12 \pm 0.05 ^f	1.49 \pm 0.19 ^d	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G2O29 (B) <i>Bacillus thuringiensis</i>	1.32 \pm 0.10 ^e	1.48 \pm 0.33 ^d	1.82 \pm 0.21 ^c	1.66 \pm 0.24 ^b	1.20 \pm 0.10 ^b	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G3O15 (B) <i>Bacillus thuringiensis</i>	1.25 \pm 0.07 ^e	1.60 \pm 0.28 ^c	1.89 \pm 0.16 ^c	1.60 \pm 0.16 ^c	1.22 \pm 0.08 ^b	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G8O49 (B) <i>Bacillus</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G8O51 (B) <i>Cellulomonas hominis</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	1.74 \pm 0.37 ^b	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G1W1 (B) <i>Micrococcus luteus</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	2.48 \pm 0.26 ^c
G1W13 (B) <i>Aureimonas frigidaquae</i>	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G3W20 (B) <i>Microbacterium</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G5W17 (B) <i>Microbacterium</i> sp	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f
G6W23 (B) <i>Pseudomonas stutzeri</i>	1.20 \pm 0.06 ^f	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	2.01 \pm 0.28 ^b
Positive control**	3.34 \pm 0.45 ^b	3.00 \pm 0.00 ^e	3.42 \pm 0.15 ^a	-	-	-	-
Negative control***	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^e	0.00 \pm 0.00 ^f

*Means followed by the same letter in the column do not differ by the Scott-Knott test ($P < 0.05$). **Positive control directly applied to the solid medium. Commercial enzymes: Protease - *Aspergillus oryzae*, α -amylase porcine pancreatic, Pectinase - *Aspergillus niger* (Sigma). ***Negative control directly applied in the solid medium. (1%) Saline solution.

ANOVA demonstrated significant differences for the enzymatic activity of protease (Table 6), where the EI ranged from 1.20 for G6W23 (B) *P. stutzeri* to 4.80 for G2O39 (A) *Staphylococcus* sp. Twelve of the endophytic bacterial isolates had an EI above 2.

In the case of amylase, there was a statistical difference between endophytes with regard to enzyme production, especially for isolate G2O43 (A) (*P. stutzeri*) with an index of 2.15. Pectin indices ranged from 1.12 to 2.32, with the best noted for the endophytes G1O38 (B) *Bacillus thuringiensis* (EI = 2.21), G3O2 (B) *Bacillus* sp (EI = 2.26), G2O43 (A)

Pseudomonas stutzeri (EI = 2.30), and G2O26 (B) *Bacillus* sp (EI = 2.26).

Enzymatic variation for lipase (Table 6) ranged from 1.24 for G1W28 (A) (*Bacillus* sp) to 2.14 for G3O25 (B) (*Acinetobacter johnsonii*), which showed the highest index for esterase (EI = 1.62). G2O29 (B) (*Bacillus thuringiensis*) had the lowest index to esterase (EI = 1.20). To cellulase, the higher EI was obtained by isolate G5O18 (B) (*Microbacterium testaceum*) (EI = 1.94). Four isolates show positive results for cellulase.

Similar to cellulase, the enzymatic activity of xylanase from endophytes also revealed positive results in four isolates, with indices between 1.52 and 2.48. The latter was found for the isolate G1W1 (B) (*Micrococcus luteus*).

These results showed that the isolate G2O39 (A) *Staphylococcus* sp possessed the greatest EI of all the tested isolates (Figure 3), with an EI of 4.80 for protease. This was significantly higher than that observed for the positive control commercial enzyme Protease - *Aspergillus oryzae* (EI = 3.34). In the case of other commercial enzymes used as positive controls (α -amylase porcine pancreatic enzyme and pectinase from *Aspergillus niger*), the EI ranging between 3.00 and 3.42, respectively, were statistically higher than endophytic bacteria tested.




Figure 3. Proteolytic activity of isolates from *Malpighia emarginata*: **a.** protease extracted from *Aspergillus oryzae* (positive control); **b.** G2O39 (A) *Staphylococcus* sp.

DISCUSSION

Sampled leaf fragments had low colonization frequency, with indices ranging from 10.4 to 21.2%, which may be due to several factors such as climate, rainfall, temperature, humidity, and the age of plants. In fact, Arnold and Herre (2003) noted that these factors affect the occurrence of endophytes. After isolation, many endophytes could not be cultivated, perhaps due to their need of culture media or specific conditions for growth (Lacava et al., 2006).

In their research on the isolation and identification of endophytic bacteria of cassava from three Brazilian States (São Paulo, Amazonas, and Bahia), Teixeira et al., (2007) listed 27 genera, with the most frequent being *Bacillus*, *Burkholderia*, *Enterobacter*, *Escherichia*, *Salmonella*, *Stenotrophomonas*, and *Serratia*.

West et al. (2010) isolated endophytic bacteria from several tissues of grapevine including the stem, root, and leaves. Most endophytes were isolated from roots, with the genus *Bacillus* spp (26% of total) being the most frequently identified strain, as well as the most frequent genus found in leaves. *B. cereus*, *B. lentimorbus*, *B. thuringiensis*, *B. megaterium*, and *B. casei* were among the species isolated, as were species of the genera *Pseudomonas* spp, *Curtobacterium* spp, *Streptomyces* spp, as well as *Enterobacter cloacae*.

Seo et al. (2010) isolated endophytic bacteria from radish leaves and roots. Leaf samples produced 140 isolates from 14 species, including *Enterobacter* sp, *B. subtilis*, *B. licheniformis*, *Stenotrophomonas maltophilia*, *Pseudomonas* sp, *Myroides odoratimimus*, *Microbacterium* sp, *Citrobacter freundii*, and others.

The taxonomic identity of several endophytes isolated from *M. emarginata* was based on molecular phylogenetic analysis by comparing ITS sequences of endophytes isolated with those deposited in GenBank, also considering the type strain sequences (when present in LPSN databases), by BLASTn. Comparison with type strain sequences provided identity rates above 90%, except for those verified for isolates G2W4 (B) and G8051 (B) (Table 2).

Among the endophytic bacteria from *M. emarginata* leaves, the presence of several genera was consistent with the results of previous studies. In most studies, the most abundant group was found to be *Bacillus*, similar to the results of the current study, in which 21 isolates were identified. These formed the most populous clade, accounting for 36.20% of the endophytes under analysis, with 19.04% identified as *B. licheniformis*, an endophyte species isolated from the leaves and roots of radish by Seo et al. (2010), and 19.04% identified as *B. thuringiensis*, an endophyte species in grape isolates reported by West et al. (2010) in isolates of banana leaves registered by Souza et al. (2014).

The genus *Staphylococcus*, which ranked second in the number of acerola isolates, was also isolated from potato. Its positive activity was also reported in an assay for growth enhancement in plants (Sessitsch et al., 2004).

Lacava et al. (2006) characterized the endophytic bacterial community of citrus and investigated its relationship with the occurrence of citrus variegated chlorosis (CVC). In asymptomatic plants, the endophyte community has been reported to be composed of the genera *Klebsiella* sp, *Acinetobacter* sp, *Curtobacterium* sp, *Pseudomonas* sp, *Enterobacter* sp, *Bacillus* spp, and *Methylobacterium mesophilicum*, whereas CVC-affected plants are preferential hosts of *Methylobacterium* spp. The genera *Pseudomonas* sp and *Enterobacter* sp were also isolated as endophytes from sugarcane (Magnani et al., 2013).

Two species of *Microbacterium*, namely, *M. testaceum* and *M. paraoxydans*, and the isolate *Cellulomonas hominis* were identified in endophytic isolates from *M. emarginata*. Within the eight genera identified from papaya tree, Thomas et al. (2007) reported the *Enterobacter* genus as usually isolated in endophyte conditions.

There are no reports in the literature on the genus *Aureimonas* isolated as endophytes, similar to that observed in the present study. Denner et al. (2003) were the first to describe the genus *Aureimonas* that, together with *Fulvimarina* and *Martella*, constitute the family Aurantimonadaceae within the order Rhizobiales (Kim et al., 2008). The genera described so far derive from corals, such as that originally reported by Denner et al. (2003) as *Aurantimonas coralicida*, from underground environments, *Aurantimonas altamirensis* (Jurado et al., 2006) and *Aurantimonas ureilytica* (Weon et al., 2007), and *Aurantimonas frigidaquae* (Kim et al., 2008) from cooling systems. *Aurantimonas coralicida* and *Aurantimonas altamirensis* were

isolated from a diseased coral (Denner et al., 2003) and from underground soil (Jurado et al., 2006). In 2008, Kim et al. studied cooling water systems in South Korea and isolated a yellow bacterial strain identified as a new species of the genus *Aurantimonas*, called *Aurantimonas frigidaquae*. This is the first report of *Aureimonas frigidaquae* as a plant endophyte. However, based on the presence or absence of a glycolipid, the genus was re-classified by Rath sack et al. (2011). Those authors suggested that the genus *Aurantimonas* could be divided, and that the species *Aurantimonas altamirensis*, *Aurantimonas ureilytica*, and *Aurantimonas frigidaquae* could be transferred to a new genus named *Aureimonas*.

The taxonomic identification of *Pseudomonas putida* was confirmed in four isolates of *M. emarginata*. Endophytic bacteria of the genus *Pseudomonas* have already been isolated from several plant species, including *Eucalyptus* sp, sugar cane, and ginger (Procópio et al., 2009; Magnani et al., 2013; Chen et al., 2014). Kumar et al. (2014) isolated endophytic bacteria from *Cassia tora* roots and identified the species *Pseudomonas putida* among the isolates.

Consistent with the results of the present study, Barretti et al. (2009) isolated *Acinetobacter johnsonii* among endophytes from the tomato plant and showed this species to cause a significant decrease in the severity of the bacterial speck caused by *Pseudomonas syringae*. The genus *Acinetobacter* was also isolated as an endophyte in studies by Procópio et al. (2009).

Most of the isolated endophytes belong to the variety Olivier when the distribution of endophytes per variety of host species and isolation temperature are taken into account. The number of endophytes belonging to the genus *Bacillus*, identified in this variety, was similar (10) for each temperature evaluated (28° and 37°C). For the genera *Staphylococcus*, *Cellulomonas*, and *Enterobacter*, an equilibrium was reported between the number of isolates of the two varieties and the temperature. The number of endophytes belonging to *Acinetobacter* and *Pseudomonas* was greatest in the variety Olivier, whereas the variety Waldy Cati 30 only hosted isolates of the genera *Micrococcus* and *Aureimonas*.

In the current analysis, 92.30% of bacteria demonstrated antagonistic activity against phytopathogenic fungi. Furthermore, 12.82% inhibited the growth of all fungi analyzed, especially the endophyte G8O43 (A) (*Bacillus* sp), which had the highest inhibition index for four of the seven phytopathogens studied (*Glomerella* sp, *F. solani*, *Alternaria* sp, and *F. oxysporum*). However, 7.69% did not have any anti-fungus activity.

Tonelli et al. (2010) investigated endophytic and epiphytic bacteria isolated from *Arachis hypogaea* L. (peanut) and tested their anti-fungal potential against four main pathogens, namely, *Sclerotinia sclerotiorum*, *S. minor*, *Sclerotium rolfsii*, and *Fusarium solani*. The results showed that 19 endophytic isolates inhibited the growth of one or more fungi when analyzed on plates with yeast extract and mannitol agar (YEMI) medium, whereas nine isolates showed an antagonistic effect with PDA medium.

In their analysis of endophytic bacteria isolated from *Theobroma cacao* leaves, Melnick et al. (2011) reported that 30.7% of the 69 isolates tested for *in vitro* antagonistic activity inhibited the growth of *Phytophthora capsici*, 28.6% inhibited *Moniliophthora roreri*, and 38.3% acted against *M. perniciosa*. The three pathogens were inhibited by 21.7% of isolates, with special reference to the species *Bacillus amyloquefaciens*, in which endophytic bacterial isolates of all species inhibited the growth of all pathogens. Similar results occurred for 80% of isolates from the species *B. subtilis*. Furthermore, Zhou et al. (2014) evaluated the antagonistic activity of *Pseudomonas fluorescens* against the fungus *Athelia rolfsii* and concluded that fungal growth was totally inhibited by endophytic bacteria.

The results of the present study showed that 64.70% of the endophytic bacteria associated with *M. emarginata* produce at least one of the evaluated enzymes. The isolate G2O26 (B) (*Bacillus thuringiensis*) generated positive results for all enzymes under analysis and should be highlighted. The endophytic bacterium G3O2 (B) (*Bacillus thuringiensis*) also provided a good result in terms of enzyme production, except in the production of esterase. Although several isolates had EIs of over 2.0, 32.45% of the 34 endophytic bacterial isolates did not possess enzymatic activity. It should also be highlighted that the isolates G1O47 (B) (*Staphylococcus* sp) and G2O39 (A) (*Staphylococcus* sp) produced proteases with higher proteolytic activity than the commercial enzyme employed, highlighting the biotechnological potential of these isolates for industries (e.g., food and leather).

According to Sivaramakrishnan et al. (2006), bacteria of the genus *Bacillus* are used widely in the production of thermostable α -amylase, which complies with industrial demands. *B. subtilis*, *B. stearothermophilus*, *B. licheniformis*, and *B. amyloliquefaciens* produce the highest amounts of this enzyme for commercial purposes and for several other applications.

Rodarte et al. (2011) assessed the capacity of microorganisms isolated from *Coffea arabica* L. grains to secrete extracellular proteases. Forty bacterial isolates were identified among the samples analyzed, of which 52.5% possessed enzymatic activity. In the present study, enzyme production in media containing hydrolyzed casein was divided into four levels, according to the diameter (cm) of the degradation halo as follows: (-) no degradation; (+) halos ranging between 0.1 and 0.5 cm; (++) halos ranging between 0.5 and 0.8 cm; and (+++) for halos ranging between 0.8 and 1.0 cm. The isolates *Acinetobacter* sp, *Bacillus subtilis*, *Bacillus megaterium*, *Enterobacter agglomerans*, *Kurthia* sp, *Serratia plymutica*, and *Tatumella pyseos* possessed the highest enzyme activity.

Castro et al. (2014) studied two species of mangrove plants, *Rhizophora mangle* and *Avicennia nitida*, and semi-quantitatively evaluated the production of amylase, esterase, lipase, cellulase, and protease from 40 endophyte bacteria. Those authors reported that 75% of bacterial isolates possessed proteolytic activity, followed by 52.5% with lipolytic activity; 45% of the bacterial isolates revealed amylolytic activity, especially for the isolate MBA2.19 *Erwinia* sp with EI = 6.83 for lipase activity. In the case of protease, the group with the highest number of enzyme-producing isolates and the best index was MBA2.29 *Stenotrophomonas* sp (EI = 2.53).

The findings of this study demonstrated that *M. emarginata* is colonized by several different species of endophytic bacteria, and nine different genera are represented within the 58 identified. The genus *Aureimonas*, represented by the species *Aureimonas frigidaquae*, is highlighted for the first time in specialized literature as an endophytic isolate. Regarding the biotechnological potential, these results show that endophytic bacteria associated with acerola are capable of inhibiting the growth of *S. sclerotiorum* (a fungus highly difficult to control by *in vitro* microorganisms) and *F. oxysporum* (fusariosis-causing pathogen in acerola). The endophytes G1O47 (B) *Staphylococcus* sp and G2O39 (A) *Staphylococcus* sp produced proteases with statistically higher activity than those obtained commercially, which increases the interest in these endophytic bacteria, due to the fact that enzyme production by microorganisms is highly profitable, reinforcing the importance of prospecting enzymes of industrial interest associated to acerola.

Conflicts of interest

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank CNPq (#311534/2014-7; #447265/2014-8) and Fundação Araucária - FA (#276/2014) for financial support and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) by the PNPd-UEM postdoctoral scholarship.

REFERENCES

- Anand D, Padmanabhan SA, Deventhiran M and Saravanan P (2014). Isolation of *Pseudomonas* sp from *Wrightia tinctoria* leaf and its 16s rRNA study. *Ind. J. Sci. Res.* 9: 28-31.
- Arnold AE and Herre EA (2003). Canopy cover and leaf age affect colonization by tropical fungal endophytes: Ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia* 95: 388-398. <http://dx.doi.org/10.2307/3761880>
- Azevedo JL, Maccheroni WJ, Pereira JO and Araujo WJ (2000). Endophytic microorganisms: a review on insect control and recent advances on tropical plants. *Electron. J. Biotechnol.* 3: 40-65. <http://dx.doi.org/10.2225/vol3-issue1-fulltext-4>
- Barretti PB, Romeiro RS, Mizubuti ESG and Souza JT (2009). Seleção de bactérias endofíticas de tomateiro como potenciais agentes de biocontrole e de promoção de crescimento. *Cienc. Agrotec.* 33: 2038-2044. <http://dx.doi.org/10.1590/S1413-70542009000700057>
- Carrington CMS and King RAG (2002). Fruit development and ripening in Barbados cherry, *Malpighia emarginata* DC. *Sci. Hortic. (Amsterdam)* 92: 1-7. [http://dx.doi.org/10.1016/S0304-4238\(01\)00268-0](http://dx.doi.org/10.1016/S0304-4238(01)00268-0)
- Castro RA, Quecine MC, Lacava PT, Batista BD, et al. (2014). Isolation and enzyme bioprospection of endophytic bacteria associated with plants of Brazilian mangrove ecosystem. *Springerplus* 3: 382. <http://dx.doi.org/10.1186/2193-1801-3-382>
- Cavichioli JC, Garcia MJM, Brida AL and Wilcken SRS (2014). Reaction in Barbados Cherry (*Malpighia emarginata* D.C.) to *Meloidogyne enterolobii*. *Rev. Bras. Frutic.* 36: 156-160. <http://dx.doi.org/10.1590/0100-2945-429/13>
- Chebotaev VK, Malfanova NV, Shcherbakov AV, Ahtemova GA, et al. (2015). Endophytic Bacteria in Microbial Preparations that Improve Plant Development. *Prikl. Biokhim. Mikrobiol.* 51: 271-277. <http://dx.doi.org/10.7868/S0555109915030058>
- Chen T, Chen Z, Ma GH, Du BH, et al. (2014). Diversity and potential application of endophytic bacteria in ginger. *Genet. Mol. Res.* 13: 4918-4931. <http://dx.doi.org/10.4238/2014.July.4.6>
- Denner EBM, Smith GW, Busse H-J, Schumann P, et al. (2003). *Aurantimonas corallicida* gen. nov., sp. nov., the causative agent of white plague type II on Caribbean scleractinian corals. *Int. J. Syst. Evol. Microbiol.* 53: 1115-1122. <http://dx.doi.org/10.1099/ijs.0.02359-0>
- Duza MB and Mastan SA (2013). Microbial enzymes and their applications - A Review. *Indo American J. Pharmaceut. Res* 3: 6208-6219.
- Ferreira DF (2011). Sisvar: a computer statistical analysis system. *Cienc. Agrotec.* 35: 1039-1042.
- Gujral MS, Agrawal P, Khetmalas MB and Pandey R (2013). Colonization and plant growth promotion of Sorghum seedlings by endorhizospheric *Serratia* sp. *Acta Biol. Indica* 2: 121-124.
- Jurado V, Gonzalez JM, Laiz L and Saiz-Jimenez C (2006). *Aurantimonas altamirensis* sp. nov., a member of the order Rhizobiales isolated from Altamira Cave. *Int. J. Syst. Evol. Microbiol.* 56: 2583-2585. <http://dx.doi.org/10.1099/ijs.0.64397-0>
- Kim MS, Hoa KTQ, Baik KS, Park SC, et al. (2008). *Aurantimonas frigidaquae* sp. nov., isolated from a water-cooling system. *Int. J. Syst. Evol. Microbiol.* 58: 1142-1146. <http://dx.doi.org/10.1099/ijs.0.65421-0>
- Kumar V, Kumar A, Pandey KD and Roy BK (2014). Isolation and characterization of bacterial endophytes from the roots of *Cassia tora* L. *Ann. Microbiol.* 65: 1391-1399. <http://dx.doi.org/10.1007/s13213-014-0977-x>
- Kusari S, Hertweck C and Spiteller M (2012). Chemical ecology of endophytic fungi: origins of secondary metabolites. *Chem. Biol.* 19: 792-798. <http://dx.doi.org/10.1016/j.chembiol.2012.06.004>
- Lacava PT and Azevedo JL (2013). Endophytic bacteria: A biotechnological potential in agrobiological system. In: *Bacteria in Agrobiological: Crop Productivity* (Maheshwari, DK, Saraf M and Aeron A, eds.). Springer, Berlin, Heidelberg, 1-44.
- Lacava PT, Andreote FD, Araújo WL and Azevedo JL (2006). Caracterização da comunidade bacteriana endofítica de citros por isolamento, PCR específico e DGGE. *Pesquisa Agropecu. Bras.* 41: 637-642. <http://dx.doi.org/10.1590/S0100-204X2006000400013>
- Magnani GS, Cruz LM, Weber H, Bessalho JC, et al. (2013). Culture-independent analysis of endophytic bacterial

- communities associated with Brazilian sugarcane. *Genet. Mol. Res.* 12: 4549-4558. <http://dx.doi.org/10.4238/2013.October.15.3>
- Melnick RL, Suárez C, Bailey BA and Backman PA (2011). Isolation of endophytic endospore-forming bacteria from *Theobroma cacao* as potential biological control agents of cacao diseases. *Biol. Control* 57: 236-245. <http://dx.doi.org/10.1016/j.biocontrol.2011.03.005>
- Nogueira CAM, Momesso CAS, Machado RLD, Almeida MTG, et al. (2004). Desempenho de Kits comerciais e protocolos laboratoriais para a extração de DNA genômico bacteriano. *Rev. Panam. Infectol.* 6: 35-38.
- Oliveira NA, Oliveira LA, Andrade JS and Chagas Júnior AF (2006). Atividade enzimática de isolados de rizóbia nativos da Amazônia central crescendo em diferentes níveis de acidez. *Ciênc. Tecnol. Aliment.* 26: 204-210. <http://dx.doi.org/10.1590/S0101-20612006000100032>
- Procópio REL, Araújo WL, Maccheroni W, Jr. and Azevedo JL (2009). Characterization of an endophytic bacterial community associated with *Eucalyptus* spp. *Genet. Mol. Res.* 8: 1408-1422. <http://dx.doi.org/10.4238/vol8-4gmr691>
- Rathsack K, Reitner J, Stackebrandt E and Tindall BJ (2011). Reclassification of *Aurantimonas altamirensis* (Jurado et al. 2006), *Aurantimonas ureilytica* (Weon et al. 2007) and *Aurantimonas frigidaquae* (Kim et al. 2008) as members of a new genus, *Aureimonas* gen. nov., as *Aureimonas altamirensis* gen. nov., comb. nov., *Aureimonas ureilytica* comb. nov. and *Aureimonas frigidaquae* comb. nov., and emended descriptions of the genera *Aurantimonas* and *Fulvimarina*. *Int. J. Syst. Evol. Microbiol.* 61: 2722-2728. <http://dx.doi.org/10.1099/ijs.0.027029-0>
- Rodarte MP, Dias DR, Vilela DM and Schwan RF (2011). Proteolytic activities of bacteria, yeasts and filamentous fungi isolated from coffee fruit (*Coffea arabica* L.). *Acta Sci. Agron.* 33: 457-464.
- Santos SML, De Vasconcelos AM, Oliveira VO, Clemente E, et al. (2012). Evaluation of Physical and Physicochemical Characteristics of *Malpighia emarginata* DC from the State of Ceará. *Int. J. Biochem. Res. Rev* 2: 152-163. <http://dx.doi.org/10.9734/IJBCCR/2012/1945>
- Schlaberg R, Simmon KE and Fisher MA (2012). A systematic approach for discovering novel, clinically relevant bacteria. *Emerg. Infect. Dis.* 18: 422-430. <http://dx.doi.org/10.3201/eid1803.111481>
- Seo WT, Lim WJ, Kim EJ, Yun HD, et al. (2010). Endophytic bacterial diversity in the young radish and their antimicrobial activity against pathogens. *J. Korean Soc. Appl. Bi.* 53: 493-503. <http://dx.doi.org/10.3839/jksabc.2010.075>
- Sessitsch A, Reiter B and Berg G (2004). Endophytic bacterial communities of field-grown potato plants and their plant-growth-promoting and antagonistic abilities. *Can. J. Microbiol.* 50: 239-249. <http://dx.doi.org/10.1139/w03-118>
- Sivaramakrishnan S, Gangadharan D, Nampoothiri KM, Soccol CR, et al. (2006). α -Amylases from microbial sources - an overview on recent developments. *Food Technol. Biotechnol.* 44: 173-184.
- Soria S, Alonso R and Bettucci L (2012). Endophytic bacteria from *Pinus taeda* L. as biocontrol agents of *Fusarium circinatum* Nirenberg & O'Donnell. *Chil. J. Agric. Res.* 72: 281-284. <http://dx.doi.org/10.4067/S0718-58392012000200018>
- Souza A, Cruz JC, Sousa NR, Procópio ARL, et al. (2014). Endophytic bacteria from banana cultivars and their antifungal activity. *Genet. Mol. Res.* 13: 8661-8670. <http://dx.doi.org/10.4238/2014.October.27.6>
- Tamura K, Stecher G, Peterson D, Filipski A, et al. (2013). MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30: 2725-2729. <http://dx.doi.org/10.1093/molbev/mst197>
- Teixeira MA, Melo IS, Vieira RF, Costa FEC, et al. (2007). Microrganismos endofíticos de mandioca de áreas comerciais e etnovarietades em três estados brasileiros. *Pesquisa Agropecu. Bras.* 42: 43-49. <http://dx.doi.org/10.1590/S0100-204X2007000100006>
- Thomas P, Kumari S, Swarna GK and Gowda TKS (2007). Papaya shoot tip associated endophytic bacteria isolated from *in vitro* cultures and host-endophyte interaction *in vitro* and *in vivo*. *Can. J. Microbiol.* 53: 380-390. <http://dx.doi.org/10.1139/W06-141>
- Tonelli ML, Taurian T, Ibáñez F, Angelini J, et al. (2010). Selection and *in vitro* characterization of biocontrol agents with potential to protect peanut plants against fungal pathogens. *J. Plant Pathol.* 92: 73-82.
- Weon HY, Kim BY, Yoo SH, Joa JH, et al. (2007). *Aurantimonas ureilytica* sp. nov., isolated from an air sample. *Int. J. Syst. Evol. Microbiol.* 57: 1717-1720. <http://dx.doi.org/10.1099/ijs.0.65035-0>
- West ER, Cother EJ, Steel CC and Ash GJ (2010). The characterization and diversity of bacterial endophytes of grapevine. *Can. J. Microbiol.* 56: 209-216. <http://dx.doi.org/10.1139/W10-004>
- Zhou JY, Zhao XY and Dai CC (2014). Antagonistic mechanisms of endophytic *Pseudomonas fluorescens* against *Athelia rolfsii*. *J. Appl. Microbiol.* 117: 1144-1158. <http://dx.doi.org/10.1111/jam.12586>